

MAIN PANEL

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**3Sixty5**

**TURF FUNGICIDE**

**ACTIVE CONSTITUENT: 365 g/L IPRADIONE**

**GROUP B FUNGICIDE**

**5 L**

A suspo-emulsion formulation, with improved rainfastness (over other water based formulations), for the control of fungal diseases in recreational turf as specified in the directions for use.

Read the attached booklet before use.

(label code)

REAR PANEL Attached to Container.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT expose product to low temperatures. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site.

If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with the eyes. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and water-proof gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre (ph: 131126).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 59278/5L/1205



BAR  
CODE

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

Bayer Environmental Science  
A Business Group of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)

Batch Number:

Date of Manufacture:

(label code)

**DIRECTIONS FOR USE:**

**PRECAUTIONS: DO NOT GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY**

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Recreational Turf	Brown Patch ( <i>Rhizoctonia solani</i> )	80 to 125 mL /10 to 150 L water /100 m <sup>2</sup>	<u>Curative Program:</u> Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. <u>For preventative treatment:</u> Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than 2 consecutive sprays of 3sixty5 or related dicarboximide fungicides unless tank mixed with a fungicide from a different activity group.
	Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	80 mL/10 to 110 L of water/100 m <sup>2</sup>	<u>For Curative Treatment:</u> Repeat treatment in 14 days. <u>For Preventative Treatment:</u> Apply monthly. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of 3sixty5 or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Winter Fusarium or Fusarium Patch ( <i>Fusarium nivale</i> ) (NOT Qld)	125 mL/10 to 150 L of water/100 m <sup>2</sup>	
	Helminthosporium disease affecting leaf, crown and roots: (Black and white <i>Helminthosporium</i> spot <i>Bipolaris</i> spp. <i>Drechslera</i> spp. <i>Exserohilum</i> spp.)	65 mL/10 to 150 L of water/100 m <sup>2</sup>	<u>For Curative Treatment:</u> Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass ( <i>Cynodon dactylon</i> ) only, use higher water volumes of 100 to 150L of water per 100m <sup>2</sup> . <u>For Preventative Treatment:</u> Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of 3sixty5 or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS:**

**Note:** 3sixty5 may be very unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding 3sixty5. A suitable registered buffering agent may have to be added to bring pH down below 7.

**Application:** Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

**TURF NUTRITION:** This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

Rear Panel Contd

---

**COMPATIBILITY:**

The use of a surfactant or spray oil is not recommended with 3sixty5 as it may result in damage to sensitive plants. DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage.

As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities.

**FUNGICIDE RESISTANCE STRATEGY:**

<b>GROUP</b>	<b>B</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

For fungicide resistance management 3sixty5 Fungicide is a Group B fungicide. Some naturally occurring individual fungi resistant to 3sixty5 Fungicide and other Group B fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 3sixty5 Fungicide and other Group B fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of 3sixty5 Fungicide to control resistant fungi.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:** DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Bayer Environmental Science  
A Business Group of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East Vic. 3123

Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)