

MAIN PANEL

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**Dedicate<sup>®</sup>**

**TURF FUNGICIDE**

**ACTIVE CONSTITUENT: 200 g/L TEBUCONAZOLE**  
**100 g/L TRIFLOXYSTROBIN**

**GROUP 3 11 FUNGICIDE**

**1 L**

For control of fungal diseases in recreational turf as specified in the Directions For Use.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE**

REAR PANEL attached to container

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT re-use empty containers for any other purpose.

### SAFETY DIRECTIONS

When opening the container and mixing and loading, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

### RE-ENTRY

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre (ph: 131126).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 62638/1 L/0709

Dedicate is a Registered Trademark of Bayer.

BAR  
CODE

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

Bayer Environmental Science  
A Business Operation of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East, Vic. 3123

Telephone: 1800 804 479

[www.bayeres.com.au](http://www.bayeres.com.au)

Batch Number:

Date of Manufacture:

(label code)

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**DEDICATE® TURF FUNGICIDE**

**Active constituent: 200 g/L TEBUCONAZOLE  
100g/L TRIFLOXYSTROBIN**

<b>GROUP</b>	<b>3</b>	<b>11</b>	<b>FUNGICIDE</b>
--------------	----------	-----------	------------------

**For control of fungal diseases in recreational turf as specified in the directions for use.**

**READ THIS BOOKLET BEFORE USE**

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT re-use empty containers for any other purpose.

### SAFETY DIRECTIONS

When opening the container and mixing and loading, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

### RE-ENTRY

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre (ph: 131126).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 62638/0709

LABEL BOOKLET (contd)

---

## **DIRECTIONS FOR USE:**

### **SPRAY DRIFT RESTRAINTS**

**DO NOT** apply with spray droplets smaller than a **MEDIUM** spray droplet category as defined by the ASAE S572 Standard. Users **MUST ONLY USE** nozzles classified as suitable for delivering a **MEDIUM** spray droplet category according to the nozzle manufacturer's specifications.

**DO NOT** apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

**DO NOT** apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are: **1.** date and start and finish times of application; **2.** location address and areas sprayed; **3.** full name of this product; **4.** amount used per hectare and number of hectares applied to; **5.** situation and disease; **6.** wind speed and direction during application; **7.** air temperature; **8.** nozzle brand, model and type and spray system pressure measured during application; **9.** name and address of person applying this product. (Additional record details may be required by the State or Territory where this product is used).

LABEL BOOKLET (contd)**MANDATORY NO-SPRAY ZONES**

**DO NOT** apply if there are aquatic and wetland areas, including aquacultural ponds or surface streams and rivers or, within **10 metres** downwind from the application area.

**DO NOT** apply if there are susceptible non-target plants within **15 metres**, downwind from the application area.

**DO NOT** apply if there are livestock, pasture or any land that is producing feed for livestock within **15 metres** downwind from the application area.

SITUATION	DISEASE	RATE	WATER RATE	CRITICAL COMMENTS
Turf	Anthracnose Brown Patch Winter Fusarium Helminthosporium Leptosphaerulina Curvularia	2 L/ha or 20 mL/100 m <sup>2</sup>	800 -1600 L per ha  8-16 L per 100 m <sup>2</sup>	For preventative control apply on a 21-28 day interval when conditions are favourable for disease development. For curative control apply on a 14 – 21 day interval. For brown patch control apply sufficient irrigation to wash Dedicate into the crown and upper root zone. This use is subject to an FRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	Ectotrophic Root Infecting Fungi (ERI)			Make monthly preventative applications in the three months preceding the disease season. Apply irrigation to wash Dedicate into the root zone. This use is subject to an FRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	Dollar Spot	3 L/ha		For preventative control apply on a 21-28 day interval when conditions are favourable for disease development. To control existing infections apply on a 14 – 21 day interval. This use is subject to an FRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	White helminthosporium	280 mL per bowling green	150 L per bowling green	Make monthly preventative applications from December to March. This use is subject to an FRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**DO NOT GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY**

**GENERAL INSTRUCTIONS:**

**Application:** Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

**TURF NUTRITION:** This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

**COMPATIBILITY:**

As formulations of other manufacturers' products are beyond the control of Bayer Environmental Science, all mixtures should be tested prior to mixing commercial quantities.

**FUNGICIDE RESISTANCE STRATEGY:****GROUP 3 | 11 FUNGICIDE**

For fungicide resistance management Dedicate is a Group 3 & 11 fungicide. Some naturally occurring individual fungi resistant to Dedicate and other Group 3 & 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Dedicate and other Group 3 & 11 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of Dedicate to control resistant fungi.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:** DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Risk to non-target insects – Dedicate may have adverse effects on some non-target beneficial insects, in particular predatory bugs. This should be taken into account where Integrated Pest Management utilising these organisms is practiced.

Bayer Environmental Science  
A Business Operation of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East Vic. 3123

Telephone: 1800 804 479  
[www.bayeres.com.au](http://www.bayeres.com.au)