

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baycor[®] 300

FUNGICIDE SPRAY

Active Constituent: 300 g/L BITERTANOL
Solvent: 539 g/L N-METHYL-2-PYRROLIDONE

GROUP	C	FUNGICIDE
-------	----------	-----------

For control of rust on beans and various ornamentals, black spot on roses and powdery mildew on begonias, eucalyptus seedlings and roses as specified in the DIRECTIONS FOR USE table

GENERAL INSTRUCTIONS

Note on ornamentals: Baycor 300 Fungicide Spray has been used on a wide range of ornamental plant species without damage. However some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions, it is advisable to treat only a small number of plants first, in order to ascertain their reaction, before treating the whole crop.

Warning: Some cultivars of Shasta daisy may be damaged, therefore before treating, test Baycor 300 on single plants.

Mixing

Mix the required quantity of Baycor 300 with water in the spray tank and agitate thoroughly. A non-ionic wetting agent may be added, if required, to ensure effective wetting of foliage. Baycor 300 is a highly effective fungicide with a broad spectrum of activity. The active constituent penetrates into plant tissue and thus has some measure of depth action. Besides displaying good protective activity, Baycor 300 has curative action on the disease.

Fungicide Resistance Warning

Baycor 300 Fungicide Spray is a member of the DMI group of fungicides. For fungicide resistance management Baycor 300 Fungicide Spray is a Group **C** fungicide. Some naturally occurring fungal populations resistant to Baycor 300 Fungicide Spray and other Group **C** fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungal will not be controlled by Baycor 300 Fungicide Spray and other Group **C** fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of Baycor 300 Fungicide Spray to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Harmful to fish. DO NOT contaminate ponds, waterways and drains with this product or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with eyes and skin. Will irritate the eyes, nose, throat and skin. Will damage eyes. Do NOT inhale vapour. When using the product wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length butyl rubber gloves, face shield and goggles. If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from www.bayercropscience.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Baycor® is a registered trademark of Bayer

APVMA Approval No.: 30381/1103

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Beans (<i>Phaseolus vulgaris</i>)	Rust	500 mL/ha	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 14 days later. A third application may be necessary where infection occurs early or disease pressure is high. Ensure thorough coverage of all foliage.
Calendula, Carnation, Chrysanthemum, Pelargonium, Snapdragon, Fuchsia, Boronia		170 mL/100 L	Use only high volume application equipment. Do NOT use misting machines. Curative: For existing infections, spray to run-off ensuring thorough coverage of all foliage including the underside of leaves in a series of 3 applications at 5 day intervals. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 to 14 days according to infection pressure.
Rose	Rust (Qld only) Black spot		Use only high volume application equipment. Do NOT use misting machines. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 to 14 days according to infection pressure or beginning immediately after onset of the disease.
Begonia, Eucalyptus seedlings	Powdery mildew	130 mL/100 L	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

Beans: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION