

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Euparen[®] Multi

500 WG FUNGICIDE

Active Constituent: 500 g/kg TOLYLFLUANID

GROUP X FUNGICIDE

For the control of black spot and grey mould and suppression of powdery mildew on strawberries

GENERAL INSTRUCTIONS

Fungicide Resistance Warning

For fungicide resistance management the product is a Group X fungicide. Some naturally occurring fungal populations resistant to the product and other Group X fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group X fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

Mixing

Add the required quantity of Euparen Multi 500 WG Fungicide by **slowly** sprinkling granules directly to the water in the spray tank while stirring or with agitators in motion.

Compatibility

Euparen Multi is compatible with most commonly used wettable powder formulations of insecticides and fungicides. Do not mix Euparen Multi with emulsifiable concentrates, or materials of a strongly alkaline nature. Do not add wetting or sticking agents to Euparen Multi sprays. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Bayer CropScience Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer CropScience cannot be responsible for the behaviour of such mixtures.

PRECAUTIONS

Re-entry Period

Do not allow entry into treated area until spray has dried. When prior entry is necessary, wear cotton overalls button to the neck and wrist and a washable hat and chemical resistant gloves.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is very highly toxic to fish and aquatic invertebrates, and moderately toxic to algae. Do not contaminate streams, rivers or waterways with the chemical or used container.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent crops, crop lands, pastures or waterways.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Puncture and bury empty container in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty container and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if inhaled. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale dust. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from www.bayercropscience.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Euparen® is a registered trademark of Bayer

NRA Approval Number 50627/0703

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE**Restraint**

DO NOT spray Euparen Multi in temperatures above 30°C.

CROP	DISEASE	RATE	APPLICATION	CRITICAL COMMENTS
Strawberries	Black spot Grey mould Powdery mildew (suppression only)	2 kg/ha	Use where spray volume is less than 1000 L/ha. Do not apply volumes less than 500 L/ha	Start spraying at the commencement of flowering and continue until the end of harvest. Apply at 7 to 10 day intervals. Use the shorter interval when conditions favour development of the disease or rapid fruit growth. When applied on a schedule Euparen Multi will suppress powdery mildew.
		200 g/100 L	Use where spray volume exceeds 1000 L/ha	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

Strawberries: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION